



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Book

Search **Nucleotide**


for

**Go****Clear**

Limits

Preview/Index

History

Clipboard

Details

**Display**

default

Show: 20

**Send to**

File

**Get Subsequence****1: M69043. Homo sapiens MAD...[gi:187290]**

Links

**LOCUS** HUMMAD3A 1550 bp mRNA linear PRI 07-MAR-1994  
**DEFINITION** Homo sapiens MAD-3 mRNA encoding I<sub>k</sub>B-like activity, complete cds.  
**ACCESSION** M69043  
**VERSION** M69043.1 GI:187290  
**KEYWORDS**  
**SOURCE** Homo sapiens (human)  
**ORGANISM** Homo sapiens  
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;  
Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.  
**REFERENCE** 1 (bases 1 to 1550)  
**AUTHORS** Haskill,S., Beg,A.A., Tompkins,S.M., Morris,J.S., Yurochko,A.D.,  
Sampson-Johannes,A., Mondal,K., Ralph,P. and Baldwin,A.S. Jr.  
**TITLE** Characterization of an immediate-early gene induced in adherent  
monocytes that encodes I kappa B-like activity  
**JOURNAL** Cell 65 (7), 1281-1289 (1991)  
**MEDLINE** 91292530  
**PUBMED** 1829648  
**COMMENT** Original source text: Homo sapiens cDNA to mRNA.  
**FEATURES**  
**source** Location/Qualifiers  
1..1550  
/organism="Homo sapiens"  
/mol\_type="mRNA"  
/db\_xref="taxon:9606"  
/cell\_type="monocyte, neutrophil"  
/tissue\_lib="pcDNA I"  
**gene** 1..1550  
/gene="MAD3"  
**CDS** 95..1048  
/gene="MAD3"  
/function="I<sub>k</sub>B-like activity"  
/codon\_start=1  
/protein\_id="AAA16489.1"  
/db\_xref="GI:187291"  
/translation="MFQAAERPQEWAEGPRDGLKKERLLDDRHDSGLDSMKDEEYEQ  
MVKELQEIRLEPQEVPGRGSEPWKQQLTEDGDSFLHLAIIEEKALTMEVIRQVKGDLA  
FLNFQNNLQQTPLHLAVITNQPEIAEALLGAGCDPELRDFRGNTPLHLACEQGCLASV  
GVLTQSCTTPHLHSILKATNYNGHTCLHLASIHYGLGIVELLVSLGADVNAQEPCNGR  
TALHLAVDLQNPDLVSLLLKGADVNRTYQGYSPYQLTWGRPSTRIQQQLGQLTLEN  
LQMLPESEDEESYDTESEFTEDELPYDDCVFGGQRLTL"  
**misc\_feature** 1135..1139  
/gene="MAD3"  
/note="attta motif"  
**misc\_feature** 1346..1350  
/gene="MAD3"  
/note="attta motif"  
**misc\_feature** 1525..1529  
/gene="MAD3"

Practitioner's Docke . MPI96-031CP1DV1CPACN1M 10/052,005

## Exhibit A

/note="attta motif"

polyA\_site

1550

/gene="MAD3"

BASE COUNT 380 a 402 c 416 g 352 t

ORIGIN

1 tgccggccgtc ccgcccccca gcgcggcagc gaggaagcag cgcgccgccc gcggcccagc  
61 gcacccgcag cagcggccgc agctcggtcg cgccatgttc caggcgccg agcgccccca  
121 ggagtgggc atggagggcc cccgcgacgg gctgaagaag gagcggctac tggacgaccg  
181 ccacgacgc ggcctggact ccatgaaaga cgaggagttac gagcagatgg tcaaggagct  
241 gcaggagatc cgcctcgagc cgcaggaggt gccgcgcggc tcggagccct ggaaggcagca  
301 gctcaccgag gacggggact cgttcctgca cttggccatc atccatgaag aaaaggcact  
361 gaccatggaa gtgatccgccc aggtgaaggg agacctggct ttcctcaact tccagaacaa  
421 cctgcagcag actccactcc acttggctgt gatcaccaac cagccagaaa ttgctgaggc  
481 acttctggga gctggctgtg atcctgagct ccgagacttt cgaggaata ccccccata  
541 ccttgcctgt gagcagggt gcctggccag cgtggagtc ctgactcagt cctgcaccac  
601 cccgcacccctc cactccatcc tgaaggctac caactacaat ggccacacgt gtctacactt  
661 agcctctatc catggctacc tgggcatctgt ggagctttt gtgtccttgg gtgtgtatgt  
721 caatgcttag gagccctgta atggccggac tgcccttac ctgcagttgg acctgcaaaa  
781 tcctgacactg gtgtcactcc ttttgaagt tgggctgtat gtcaacagag ttacctacca  
841 ggcttattct ccctaccagc tcacctgggg cgcggccaaac acccggtatc agcagcagct  
901 gggccagctg acactagaaa accttcagat gctggccagag agtgaggatg aggagagcta  
961 tgacacagag tcagagttca cggagttcac agaggacgg ctggccatgt atgactgtgt  
1021 gtttggaggg cagcgtctga cgttatgagt gcaaaggggc taaaagaaca tggacttgt  
1081 tatttgtaca aaaaaaaaaat ttatttttc taaaaaaaaa aaaaaaaaaga aaaaaattaa  
1141 agggtgtact tatattccaca ctgcacactg cctagcccaa aacgtcttat tgggttagga  
1201 tcagccctca ttttggct tttgttaact tttttaggg gacgagaaag atcattgaaa  
1261 ttctgagaaa acttcttta aacctcacct ttgtgggtt tttggagaag gttatcaaaa  
1321 atttcatggg aggaccatattt tttatattt tttgtctcg agtactgac cccagtggta  
1381 tcctgtgaca tgtaacagcc aggagtgtt aacgttcagt gatgtgggtt gaaaagttac  
1441 tacctgtcaa ggtttgtgtt accctccctgt aatgggtta cataatgtat tgggttaat  
1501 tattttggta ctttatgtat gtatatttta taaagagatt ttacaaatg

//

[Disclaimer](#) | [Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)

May 20 2003 11:20:12



## Sequence Revision History

PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Books

Find (Accession, GI number or Fasta style SeqId) 

Go

Clear

About Entrez

Search for Genes  
LocusLink provides  
curated information for  
human, fruit fly, mouse,  
rat, and zebrafish

Entrez Nucleotide  
Help | FAQ

Batch Entrez: Upload a  
file of GI or accession  
numbers to retrieve  
sequences

Check sequence  
revision history

How to create WWW  
links to Entrez

LinkOut

Cubby

Related resources  
BLAST

Reference sequence  
project


Submit to GenBank

## Revision history for "M69043"

GI	Version	Update Date	Status
187290	1	Mar 7 1994 17:00	Live
187290	1	Jul 26 1993 19:32	Dead
187290	1	Apr 27 1993 12:57	Dead

Accession M69043 was first seen at NCBI on Apr 27 1993 12:57

[Disclaimer](#) | [Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Bc

Search for 

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

 1: Proc Natl Acad Sci U S A. 1993 Mar 15;90(6):2532-6.

Related Articles, Li

FREE full text article at  
[www.pnas.org](http://www.pnas.org)PubMed Central  
access FREE full text articles**Mutual regulation of the transcriptional activator NF-kappa B and its inhibitor, I kappa B-alpha.****Brown K, Park S, Kanno T, Franzoso G, Siebenlist U.**

Laboratory of Immunoregulation, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD 20892.

The NF-kappa B transcription factor complex is sequestered in the cytoplasm by the inhibitory protein I kappa B-alpha (MAD-3). Various cellular stimuli relieve this inhibition by mechanisms largely unknown, leading to NF-kappa B nuclear localization and transactivation of its target genes. It is demonstrated here with human T lymphocytes and monocytes that different stimuli, including tumor necrosis factor alpha and phorbol 12-myristate 13-acetate, cause rapid degradation of I kappa B-alpha, with concomitant activation of NF-kappa B, followed by a dramatic increase in I kappa B-alpha mRNA and protein synthesis. Transfection studies reveal that the I kappa B-alpha mRNA and the encoded protein are potently induced by NF-kappa B and by homodimers of p65 and of c-Rel. We propose a model in which NF-kappa B and I kappa B-alpha mutually regulate each other in a cycle: saturating amounts of the inhibitory I kappa B-alpha protein are destroyed upon stimulation, allowing rapid activation of NF-kappa B. Subsequently, I kappa B-alpha mRNA and protein levels are quickly induced by the activated NF-kappa B. This resurgence of I kappa B-alpha protein acts to restore an equilibrium in which NF-kappa B is again inhibited.

PMID: 8460169 [PubMed - indexed for MEDLINE]

 Display  Abstract  Show: 20  Sort  Send to  Text
[Write to the Help Desk](#)[NCBI | NLM | NIH](#)[Department of Health & Human Services](#)[Freedom of Information Act | Disclaimer](#)